Rheumatology Infusion Enrollment Form R-Z

Patient Data	Patient Name: SSN #: XXX-XX- Address: Home Phone:Cell Alternate Caregiver Name:	Phone:	Sex: AMale Female Known Allergies: City: Primary Language: Phone of Caregiver:	State:	Zip:		
lns. Data	Primary Insurance: Policy:		Secondary Insurance: Policy:				
rmation	To expedite prior authorization, please attach PPD results, lab results, clinical office notes, and past treatment history. Diagnosis : M06.9 Rheumatoid Arthritis M31.3 Granulomatosis with Polyangiitis M31.7 Microscopic Polyangiitis M32.9 Systemic Lupus M32.14 Lupus Nephritis L40.52 Psoriatic Arthritis M45.9 Ankylosing Spondylitis Other:						

Info	Does patient have a latex allergy? \Box Yes \Box No	Has patient been test	ed for Hepatitis B? \Box Yes	□No	If positive, has treatment been initiated? Types	No
cal	Has patient received PPD (tuberculosis) skin tes	it: 🛛 Yes 🔍 No 🛛 Date:			Results: 🗖 Positive 📮 Negative	
Clini	Has patient previously been treated for this cor	ndition: 🛛 Yes 🔍 No	Joints Affected:			

If yes, medication/therapy failed (length of therapy): __

SPECIALTY PHARMACY

1							
	Medication	Strength	Directions	Quantity	Refills		
	 Remicade[®] Avsola[™] infliximab biosimilar Inflectra[®] infliximab biosimilar Renflexis infliximab biosimilar 	100mg SDV vial	Rheumatoid Arthritis Induction: Infuse 3mg/kg intravenously at weeks 0, 2, and 6	Quantity Sufficient x3 doses	0 refills		
			Aaintenance: Infuse 3mg/kg intravenously every 8 weeks Patient weightkg	Quantity Sufficient for 8 weeks			
			Ankylosing Spondylitis Induction: Infuse 5mg/kg intravenously at weeks 0, 2, and 6 	Quantity Sufficient x3 doses	0 refills		
			Aaintenance: Infuse 5mg/kg intravenously every 6 weeks Patient weightkg	Quantity Sufficient for 6 weeks			
			Psoriatic Arthritis Induction: Infuse 5mg/kg intravenously at weeks 0, 2, and 6	Quantity Sufficient x3 doses	0 refills		
R		Quantity Sufficient for 8 weeks					
Prescription	 Rituxan[®] Ruxience rituximab biosimilar Truxima[®] rituximab biosimilar 	500mg/50ml vial	Rheumatoid Arthritis Infuse two 1000mg doses intravenously 2 weeks apart. Repeat every weeks (no sooner than every 16 weeks)	#4 vials			
			Granulomatosis with Polyangiitis & Miscroscopic Polyangiitis (Rituxan & Truxima ONLY) □ Induction: Infuse 375mg/m ² intravenously once weekly for 4 weeks Patient weightkg Patient heightinches	Quantity Sufficient x4 doses	0 refills		
			□ First maintenance dose: Infuse two 500mg doses intravenously 2 weeks apart, begin maintenance in 6 months	#2 vials	0 refills		
			 Final maintenance dose: Infuse 500mg intravenously once every 6 months 	#1 vial			
		50mg/4ml vial	□ Induction: Infuse 2mg/kg intravenously at weeks 0 and 4	Quantity Sufficient x2 doses	0 refills		
	☐ Simponi Aria®		Maintenance: Infuse 2mg/kg intravenously every 8 weeks Patient weightkg	Quantity Sufficient for 8 weeks			

Is patient new to this therapy:	I YES I NO	Ship to	: 🗆 Office 🗆 Othe	r	Need by Date:
Prescriber Name:			DEA#:		NPI:
Practice Name:			Contact:		
Address:			Phone:		Fax:
City: State:			Key Contact#:		

My signature below authorizes Value Specialty Pharmacy staff to act as my authorized agent to complete the insurance prior authorization process for my patient listed above. My authorization shall include any required signatures by Value Specialty Pharmacists on my behalf to facilitate this process and acknowledge their authorized access to necessary healthcare data to complete said process.

Date:

fax referral to: **844-8<u>12-622</u>7**

prescriber writes "DAW" in the box to the right.

phone: 855-265-8008 | www.vsprx.com

This prescription will be filled generically unless