VALUE Crohn's/Ulcerative Colitis Enrollment Form S-Z

Patient Data	Patient Name:	Known Allergies:				
lns. Data		Secondary Insurance: Policy: Group#: apply for manufacturer copay card. Email must be provided above.				
Diagnosis	 K50.00 Crohn's Disease of small instestine without complications K50.10 Crohn's Disease of large instestine without complications K50.80 Crohn's Disease of both small and large instestine without complications K50.90 Crohn's Disease, unspecified, without complications Other: ICD 10: 	 K51.20 Ulcerative (chronic) proctitis without complications K51.30 Ulcerative (chronic) rectosigmoiditis without complications K51.50 Left sided colitis without complications K51.80 Other ulcerative colitis without complications K51.90 Ulcerative colitis, unspecified, without complications Other: ICD 10: 				
Clinical Information						

	Medication	Strength	Directions	Quantity	Refills
	Simponi [®]	 100mg/ml PFS 100mg/ml Smartject[®] 	Induction: Inject 200mg (2 injections) SQ at week 0, then inject 100mg (1 injection) SQ at week 2	#3	0 refills
	only indicated for UC	Autoinjector	Maintenance: Inject 100mg SQ every 4 weeks	#1	
		600mg/10ml vial	□ Induction: Infuse 600mg intravenously at weeks 0, 4, and 8	#3	0 refills
	Skyrizi® only indicated for Crohn's	 180mg/1.2ml cartridge 360mg/2.4ml cartridge 180mg/1.2ml On Body 360mg/2.4ml On Body 	Maintenance: Inject 1 cartridge SQ at week 12, then every 8 weeks thereafter	#1	
Prescription	□ Stelara®	130mg/26ml vial	□ Induction: (Dosed by weight) Infusemg intravenously as single dose, begin maintenance in 8 weeks Up to $55kg \rightarrow 260mg = 2$ vials Greater than $55kg$ to $85kg \rightarrow 390mg = 3$ vials Greater than $85kg \rightarrow 520mg = 4$ vials Patient weightkg	Quantity Sufficient x1 dose	0 refills
		 45mg/0.5ml vial 90mg/ml PFS 	Maintenance: Inject 90mg SQ 8 weeks after initial IV dose, then every 8 weeks thereafter	Quantity Sufficient x1 dose	
	❑ Xeljanz [®]	10mg tablet	Induction: Take 1 tablet orally twice daily for 8 weeks	#60	1 refill
	only indicated for UC	5mg tablet	Maintenance: Take 1 tablet orally twice daily	#60	
	🖵 Xeljanz XR®	22mg tablet	□ Induction: Take 1 tablet orally once daily for 8 weeks	#30	1 refill
	only indicated for UC	11mg tablet	Aaintenance: Take 1 tablet orally once daily	#30	
	Zeposia [®]	Starter Pack	Take orally per titration pack instructions	#1 pack	0 refills
	only indicated for UC	0.92mg capsule	Take 1 capsule orally once daily	#30	

Is patient new to this therapy: 🗆 YES 🗆 NO 📔 Ship to: 🗆 Patient 🗔 Office 🗔 Other 📋 Need by Date:____

Date:

Prescriber Name:	DEA#:NPI:
Practice Name:	Contact:
Address:	Phone:Fax:
City: State: Zip:	Key Contact#:

My signature below authorizes Value Specialty Pharmacy staff to act as my authorized agent to complete the insurance prior authorization process for my patient listed above. My authorization shall include any required signatures by Value Specialty Pharmacists on my behalf to facilitate this process and acknowledge their authorized access to necessary healthcare data to complete said process.

fax referral <u>to: **844**-**812-6**227</u>

e: _____ prescriber writes "DAW" in the box to the right. phone: 855-265-8008 | www

This prescription will be filled generically unless

www.vsprx.com